Zentrum für Technomathematik
Fachbereich 3 – Mathematik und Informatik

Differentiability of Consistency Functions

Christof Büskens Matthias Gerdts

Report 04–10

Berichte aus der Technomathematik

Report 04–10 August 2004

Differentiability of Consistency Functions for DAE Systems

C. Büskens
Zentrum für Technomathematik
Universität Bremen, Bremen, Germany
and
M. Gerdts
Lehrstuhl für Angewandte Mathematik
Universität Bayreuth, Bayreuth, Germany

Abstract. In the present paper, parametric initial-value problems for differential–algebraic (DAE) systems are investigated. It is known that initial values of DAE systems must satisfy not only the original equations in the system but also derivatives of these equations with respect to time. Whether or not this actually imposes additional constraints on the initial values depends on the particular problem.

Often the initial values are not determined uniquely, so that the resulting degrees of freedom can be used to optimize a given performance index. For this purpose, a class of functions is defined which will be called consistency functions. These functions map a set of parameters, which also include those undetermined initial values, to consistent initial values for the DAE system.

Because of frequent gradient evaluations of the performance index and the constraints with respect to these system parameters needed by many optimization procedures, we state conditions such that the consistency functions represent differentiable functions with respect to these parameters.

Several examples are provided to illustrate the verification of the theoretical assumptions and their differentiability properties.
Key Words. Parametric DAE systems, consistent initial values, sensitivity analysis.

1 Introduction

Many engineering and scientific problems are described by systems of *differential–algebraic equations* (DAEs). Typical applications for such DAE systems arise in multi–body dynamics, process engineering, electric circuit simulation, robot path planning problems or singular and constrained arcs in optimal control problems. Though DAE systems are easy to formulate and are close to the *'engineer's way of thinking'*, they pose several numerical problems: *DAE's are not ODE's*; see Ref. 1.

Besides ill-conditioning, stiffness and stability problems in the numerical solution of such DAE systems, the calculation of *consistent initial values* for nonlinear DAE systems is a demanding problem. Several methods are proposed in the literature. Among others, techniques are discussed that are based on index reduction methods, artificial integration steps (Ref. 2), Taylor series expansions (Ref. 3), graph theoretic algorithms (Ref. 4), projection methods (Ref. 5), or methods that set up the derivative array equations (Refs. 6-7). In all of these articles, a unique solution for the consistent initial values is assumed.

In Büskens and Gerdts (Refs. 8-9), numerical methods are discussed where consistent initial values can be calculated even in the case that no unique solution exists. The proposed methods are based on finite dimensional optimization, where the non–uniqueness is used for an optimal exploitation of the degrees of freedom left in the consistent initial values. An objective function to be defined and certain equality constraints are introduced to guarantee both consistency and optimality of the free initial values. In this article we generalize the numerical investigations of Ref. 8 to a more general class of *consistency functions*. Hereby, a consistency function will be defined as a function which maps a set of parameters or variables to a locally uniquely defined consistent initial value for a given DAE system.

Sensitivity analysis is concerned with the behavior of solutions of a given problem with respect to parameter variations or uncertainties. *Solution differentiability*, i.e., the differentiability w.r.t. these parameters, is of particular importance. The theoretical framework for a sensitivity analysis of optimization problems has been developed by Fiacco (Ref. 10). In this book, the main result on solution differentiability is based on second order sufficient conditions (SSC) and their numerical verification. Since the computation of consistency functions is phrased in terms of an optimization problem, we shall be able to derive conditions which ensure, roughly speaking, that the consistency functions become differentiable functions of special parameters. Second order sufficient conditions (SSC) are used to proof the stability of the solution and the existence of *sensitivity differentials*.

The verification of SSC and the calculation of sensitivity differentials with respect to the parameters by numerical methods, has been proven to be a very powerful and helpful instrument for a multitude of mathematical problems, see Büskens (Ref. 11). The investigation of differentiability properties of consistent initial values with respect to parameters is not only an interesting problem on its own, but also in combination with direct shooting methods for the numerical solution of DAE optimal control problems these sensitivity differentials play a substantial role, see Gerdts (Ref. 12). The intention of this paper is to give the theoretical basis for the applicability of the numerical methods mentioned before.

The paper is organized as follows: First we give a short overview on DAE systems and some of their formulations (Section 2). The index of DAE systems and the consistency of their initial values are discussed in order to define consistency functions. This leads to a nonlinear programming problem (NLP). Well-known necessary and second order sufficient conditions for NLPs are presented in Section 3 in order to establish conditions such that solutions of the NLPs are locally unique. Furthermore, strong differentiability conditions for consistency functions are formulated by

applying these results. By extracting relevant information, these conditions can be weakened as described in Section 4. Several examples are discussed in Section 5, which show the applicability of the proposed ideas. A short conclusion completes the paper.

2 Overview and Formulation of the Problem

In the following we investigate parametric DAE systems in the most general *implicit form*. Without loss of generality, we restrict the discussion to autonomous problems:

$$F(x(t), \dot{x}(t), p) = 0, \qquad \text{for all} \quad t \in [t_0, t_f], \tag{1}$$

with a sufficiently smooth function $F : \mathbb{R}^n \times \mathbb{R}^n \times P \to \mathbb{R}^n$, $P \subseteq \mathbb{R}^m$. The argument $x(t) \in \mathbb{R}^n$ denotes the state of the dynamical system (1) at time $t \in [t_0, t_f]$, $\dot{x}(t)$ its derivative w.r.t. t, while p represents an additional (fixed) parameter.

If the Jacobian $F_{\dot{x}} := \partial F / \partial \dot{x}$ is nonsingular for all $t \in [t_0, t_f]$, equation (1) describes an *implicit ordinary differential equation system* (IODE system). In the following we confine the discussion to problems with singular Jacobian $F_{\dot{x}}$ (Refs. 1,13). In this case, Eq. (1) contains differential equations as well as algebraic equations and is called a *differential-algebraic equation system*.

DAE systems arising from engineering applications often have a special structure. For instance dynamic models of chemical engineering processes and electric circuits are described by DAE systems, which are linear in $\dot{x}$:

$$M(x(t), p)\, \dot{x}(t) = f(x(t), p), \tag{2}$$

where $M(x(t), p)$ is singular. Mechanical multi–body systems can be transformed to *semi-explicit* DAE systems:

$$\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \dot{x}(t) \\ \dot{y}(t) \end{pmatrix} = \begin{pmatrix} \dot{x}(t) \\ 0 \end{pmatrix} = \begin{pmatrix} f(x(t), y(t), p) \\ g(x(t), y(t), p) \end{pmatrix} \tag{3}$$

where the matrix M in (2) is characterized by a special structure. Note that only for the differential variables $x(t) \in \mathbb{R}^{n_x}$ a differential equation $\dot{x}(t) = f(x(t), y(t), p)$ is given explicitly and that the dimensions of $y(t)$ and $g(x(t), y(t), p)$ are the same. The *algebraic variable* $y(t) \in \mathbb{R}^{n_y}$ is implicitly given by the *algebraic constraint* $0 = g(x(t), y(t), p)$ or their derivatives w.r.t. time. Therefore in contrast to ODE systems initial value problems for a semi-explicit DAE system with initial values $x(t_0) = x_0$ and $y(t_0) = y_0$ are in general not solvable for arbitrary values (x_0, y_0). This holds for problems of form (1) as well as of form (2). Since the more common formulations (2) and (3) are only special cases of (1), we restrict our considerations to the more general implicit form (1).

The *differentiation index* describes a measure how hard the calculation of consistent initial values is (Ref. 14):

Definition 2.1 (Differentiation Index and Consistency) A natural number $k \in \mathbb{N}_0$ is called the differentiation index of the DAE (1) if it is the smallest number such that from

$$\begin{aligned} F(x(t), \dot{x}(t), p) &= 0, && \text{(4a)} \\ \frac{d}{dt} F(x(t), \dot{x}(t), p) &= 0, && \text{(4b)} \\ &\vdots \\ \frac{d^k}{dt^k} F(x(t), \dot{x}(t), p) &= 0, && \text{(4c)} \end{aligned}$$

an explicit ODE system of type

$$\dot{x}(t) = \Psi(x(t), p) \tag{5}$$

can be derived by algebraic manipulations. A set of initial values $(x_0, \dot{x}_0) \in \mathbb{R}^{2n}$ for the DAE system (1) of differentiation index k at $t = t_0$ is called consistent, if $(x_0, \dot{x}_0)$ fulfills Eqs. (4) and $\dot{x}_0 = \Psi(x_0, p)$. We call the DAE (1) solvable if there exist consistent initial values such that the ODE system (5) is solvable for each consistent initial value.

The derivatives in Eq. (4) are short-hand notations. For example, it holds

$$\frac{d}{dt} F(x(t), \dot{x}(t), p) = F_x(x(t), \dot{x}(t), p) \cdot \dot{x}(t) + F_{\dot{x}}(x(t), \dot{x}(t), p) \cdot \ddot{x}(t).$$

The higher order derivatives are defined recursively, as described in more detail in Leimkuhler et al. (Ref. 3). In particular, notice that the k^{th} derivative of F includes the derivatives $x, \dot{x}, \ddot{x}, \ldots, x^{(k+1)}$. Implicitly it is assumed that these derivatives actually exist. Furthermore, notice that a consistent initial value in general depends on the parameters p.

Note that an ODE system (the Jacobian $F_{\dot{x}}$ is nonsingular) has the differentiation index $k = 0$. Depending on the index, consistent initial values x_0 yield initial values $\dot{x}_0$ for the first derivatives by equation (5) and must satisfy (4):

$$\begin{aligned} F(x_0, \dot{x}_0, p) &= 0, && \text{(6a)} \\ \frac{d}{dt} F(x_0, \dot{x}_0, p) &= 0, && \text{(6b)} \\ &\vdots \\ \frac{d^k}{dt^k} F(x_0, \dot{x}_0, p) &= 0. && \text{(6c)} \end{aligned}$$

In general, the user may have some information about the initial state. Hence the following problem can be stated:

Problem 2.1 (Consistent Initial Values) Find consistent initial values $(x(t_0), \dot{x}(t_0))$ such that the DAE system (1) can be solved subject to an additional set of equality constraints defined by the function $\varphi : \mathbb{R}^n \times \mathbb{R}^n \times P \to \mathbb{R}^r$, $r \leq n$:

$$\begin{aligned} F(x(t), \dot{x}(t), p) &= 0, \qquad \text{for all} \quad t \in [t_0, t_f], && \text{(7a)} \\ \varphi(x(t_0), \dot{x}(t_0), p) &= 0. && \text{(7b)} \end{aligned}$$

In the literature the special case $\varphi(x(t_0), \dot{x}(t_0), p) := x(t_0) - p$ often can be found.

Often, consistent initial values are determined uniquely by Eqs. (7). In this paper we are particularly interested in the case where consistent initial values are not uniquely determined for a given parameter p. For this purpose we introduce the concept of *consistency functions*.

Definition 2.2 (Consistency Function) Let $P \subseteq \mathbb{R}^m$ and $con : P \to \mathbb{R}^{2n}$ be a function with $con(p) = (con_1(p), con_2(p))$, $con_1, con_2 : P \to \mathbb{R}^n$, that solves Problem 2.1 for all $p \in P$, i.e., $F(con_1(p), con_2(p), p) = 0$ and $\varphi(con_1(p), con_2(p), p) = 0$. Then con is called a consistency function.

In the following we will concentrate on C^1-consistency functions which have a wide area of applications, e.g. in optimization problems and optimal control problems. Our intention is to define a class of consistency functions for problems with both non-unique and unique solutions. Hence our proposal is to investigate consistency functions which are defined by the solutions of nonlinear optimization problems, e.g.,

$$\begin{aligned} \text{(NLP1(p))} \qquad \min_z \quad & g(z, p), && \text{(8a)} \\ \text{s.t.} \quad & F(x(t_0), \dot{x}(t_0), p) = 0, && \text{(8b)} \\ & \frac{d}{dt} F(x(t_0), \dot{x}(t_0), p) = 0, && \text{(8c)} \\ & \vdots \\ & \frac{d^k}{dt^k} F(x(t_0), \dot{x}(t_0), p) = 0, && \text{(8d)} \\ & \varphi(x(t_0), \dot{x}(t_0), p) = 0. && \text{(8e)} \end{aligned}$$

The objective function $g(z, p)$ has to be defined here appropriately for $z := (x(t_0), \dot{x}(t_0), \ldots, x^{(k+1)}(t_0))$. Let $\bar{x}(p) = (\bar{x}_0(p), \ldots, \bar{x}_0^{(k+1)}(p))^T$ be an optimal solution of (8), then $con(p) := (\bar{x}_0(p), \dot{\bar{x}}_0(p))^T$ defines a consistency function.

In the sequel we will formulate conditions such that the consistency function defined by the solution of (8) is a C^1 function.

Please note that if no unique consistent initial value is given by the equality constraints in (8), we have to define the objective $g(z, p)$ in (8) such that the optimal solution of (NLP1(p)) results in a locally unique solution.

In the sequel we tacitly assume that problem 2.1 is solvable.

3 Strong Solution Differentiability of Consistency Functions

A first step for the solution of problem (8) is to formulate necessary conditions. Problem (8) represents a finite dimensional nonlinear optimization problem (NLP) with *equality* constraints. Hence let

$$G(z,p) \;:=\; \begin{pmatrix} F(x(t_0),\dot{x}(t_0),p) \\ \frac{d}{dt}F(x(t_0),\dot{x}(t_0),p) \\ \vdots \\ \frac{d^k}{dt^k}F(x(t_0),\dot{x}(t_0),p) \\ \varphi(x(t_0),\dot{x}(t_0),p) \end{pmatrix} \tag{9}$$

for $G : \mathbb{R}^{(k+2)n} \times P \to \mathbb{R}^{(k+1)n+r}$, and z defined as in (8). Then Problem (8) is of form below:

$$\text{(NLP(p))} \qquad \min_z \quad g(z,p), \tag{10a}$$

$$\text{s.t.} \quad G(z,p) = 0. \tag{10b}$$

Definition 3.1 (Admissible Set, Local Minimum)

(a) The set

$$S(p) := \{z \in \mathbb{R}^{(k+2)n} | G(z,p) = 0\} \tag{11}$$

is called a set of admissible variables or admissible set. A variable $z \in S(p)$ is called admissible variable.

(b) A variable $\bar{z} \in S(p)$ is called a local, resp. strong local minimum of Problem (10), if a neighborhood $V \subseteq \mathbb{R}^{(k+2)n}$ of $\bar{z}$ exists, such that

$$g(\bar{z},p) \leq g(z,p) \text{ for all } z \in S(p) \cap V$$

resp.

$$g(\bar{z},p) < g(z,p) \text{ for all } z \in S(p) \cap V, z \neq \bar{z}.$$

Let us introduce the Lagrangian function for the nonlinear optimization problem (NLP(p))

$$L \;:\; \mathbb{R}^{(k+2)n} \times \mathbb{R}^{(k+1)n+r} \times P \longrightarrow \mathbb{R}, \tag{12a}$$

$$L(z,\mu,p) \;:=\; g(z,p) + \mu^T G(z,p), \tag{12b}$$

with multiplier $\mu \in \mathbb{R}^{(k+1)n+r}$, where $(\cdot)^T$ denotes the transpose. Herewith first order necessary optimality conditions can be formulated, cf., e.g., Fletcher (Ref. 15):

Theorem 3.1 (Strong Necessary Optimality Conditions for (NLP(p))) Let g and G be continuously differentiable with respect to z in a neighborhood of a local minimum $\bar{z}$ for Problem (10). Furthermore let the Jacobian $G_z(\bar{z}, p)$ be of full rank. Then there exist a uniquely determined multiplier $\mu \in \mathbb{R}^{(k+1)n+r}$ satisfying

$$L_z(\bar{z}, \mu, p) = g_z(\bar{z}, p) + \mu^T G_z(\bar{z}, p) = 0. \tag{13}$$

Besides necessary conditions, second order sufficient conditions (SSC) have to be checked to ensure the local optimality of solutions. These SSC play an important role for selecting optimal solutions. Another important aspect of SSC appears in the *sensitivity analysis* of Problem (10) where SSC are used to proof the existence of *sensitivity differentials*. SSC for problems of form (NLP(p)) can be validated numerically using linear algebra techniques.

Fiacco (Ref. 10) has derived conditions which ensure that solutions (z, μ) of Eq. (13) become differentiable functions of the parameter p. Let us fix a *reference* or *nominal* parameter p^0 to conduct a local sensitivity analysis. Furthermore let us consider problem (NLP(p^0)) as the *unperturbed* or *nominal* problem. We assume that there exists a local solution (z^0, μ^0) of the reference problem (NLP(p^0)) satisfying the necessary KKT conditions (13) for the nominal parameter p^0.

As a first step let us summarize different assumptions to formulate conditions for the solution differentiability of the optimal solutions of (8). We focus on the vector valued function $G(z, p)$ in Eq. (9).

Assumption 3.1

(i) Let the functions $G(z, p)$ and $g(z, p)$ be twice continuously differentiable with respect to z in a neighborhood of $z_0 := (x_0, \dot{x}_0, \ldots, x_0^{(k+1)})$ for the nominal parameter $p = p^0$.

(ii) Let the rank of the Jacobian of G be maximal, i.e.,

$$\text{rank}(G_z(x_0, \dot{x}_0, \ldots, x_0^{(k+1)}, p^0)) = (k+1)n + r. \tag{14}$$

(iii) Assume that there exists a multiplier $\mu_0 \in \mathbb{R}^{(k+1)n+r}$, such that z_0 and μ_0 satisfy the necessary optimality conditions of Theorem (3.1) with the Lagrangian

$$L(z_0, \mu_0, p^0) = g(x_0, \dot{x}_0, \ldots, x_0^{(k+1)}, p^0) + \mu_0^T G(x_0, \dot{x}_0, \ldots, x_0^{(k+1)}, p^0). \tag{15}$$

(iv) Let the functions $g_z(z, p)$, $G_z(z, p)$ and $G(z, p)$ be continuously differentiable with respect to p in a neighborhood of z_0 and p^0.

(v) The Hessian of the Lagrangian is positive definite on

$$\text{Ker}(G_z(x_0, \dot{x}_0, \ldots, x_0^{(k+1)}, p^0)). \tag{16}$$

Due to the special structure of the derivatives in Eq. (4) we find that Assumption 3.1 (i) is equivalent to

Assumption 3.1 (i') Let the DAE in (1) be of index k. Moreover let the function $F(x, \dot{x}, p)$ be $k+2$ times continuously differentiable with respect to x and $\dot{x}$, let the function $\varphi(x, \dot{x}, p)$ be twice continuously differentiable with respect to x and $\dot{x}$ and let the function $g(z, p)$ be twice continuously differentiable with respect to z in a neighborhood of z_0 for the nominal parameter $p = p^0$.

Similarly, Assumption 3.1 (iv) can be expressed in terms of the components of G in (9).

Note, that if Assumptions 3.1 (i)-(iii) hold, $z_0 = (x_0, \dot{x}_0, \ldots, x_0^{(k+1)})$ and μ_0 satisfy the conditions of Theorem 3.1, especially μ_0 is unique.

Then it holds, cf. Fiacco (Ref. 10):

Theorem 3.2 (Strong Sensitivity Analysis for Solutions of (NLP1(p))) Let Assumption 3.1 be fulfilled for the NLP (8). Then,

(a) $z_0 = (x_0, \dot{x}_0, \ldots, x_0^{(k+1)})$ is a strong local minimum of (8) and fulfills Eqs. (6). In particular, $(x_0, \dot{x}_0)$ is a consistent initial value,

(b) there exists a neighborhood $P^0 \subseteq P$ of p^0 and unique continuously differentiable functions $x_0^i : P^0 \longrightarrow \mathbb{R}^n$, $i = 0, \ldots, k+1$, and $\mu_0 : P^0 \longrightarrow \mathbb{R}^{(k+1)n+r}$ with

  (i) $x_0^{(i)}(p^0) = x_0^{(i)}$, $i = 0, \ldots, k+1$,

  (ii) $\mu_0(p^0) = \mu_0$,

  (iii) $(x_0(p), \ldots, x_0^{(k+1)}(p))$, $\mu_0(p)$ satisfy the conditions in Assumption 3.1 for the perturbed problem (NLP1(p)) for all $p \in P^0$.
  In particular $(x_0(p), \ldots, x_0^{(k+1)}(p), \mu_0(p))$ is a unique strong local minimum of (NLP1(p)),

c) the first order derivatives of $(x_0(p), \ldots, x_0^{(k+1)}(p))$ and $\mu_0(p)$ are given by

$$\begin{pmatrix} \frac{dx_0}{dp}(p^0) \\ \vdots \\ \frac{dx_0^{(k+1)}}{dp}(p^0) \\ \frac{d\mu_0}{dp}(p^0) \end{pmatrix} = - \begin{pmatrix} L_{zz}(z_0, \mu_0, p^0) & G_z(z_0, p^0)^\top \\ G_z(z_0, p^0) & 0 \end{pmatrix}^{-1} \begin{pmatrix} L_{zp}(z_0, \mu_0, p^0) \\ G_p(z_0, p^0) \end{pmatrix}. \tag{17}$$

As a direct consequence of Theorem 3.2 we find:

Theorem 3.3 (Strong Differentiability Conditions for the Consistency Function con(p)) Let Assumptions 3.1 be fulfilled for the NLP (8) with reference parameter $p = p^0$. Then, there exists a neighborhood $P^0 \subseteq P$ of p^0 such that $con(p) := (x_0(p), \dot{x}_0(p))^T$ defines a C^1 consistency function on P^0.

Assumption 3.1 describes sufficient conditions needed to proof the differentiability of consistency functions defined by (8). However, we can in general not expect that all of these conditions are fulfilled as the simple Example 3.1 shows:

Example 3.1 *Let* $z := (x_1(t_0), x_2(t_0), \ldots, x_1^{(k+1)}(t_0), x_2^{(k+1)}(t_0))$ *and* $t_0 = 0$*:*

$$\min_z \quad x_2^2(t_0), \tag{18a}$$

$$s.t. \quad \dot{x}_1(t) = \begin{cases} \frac{1}{6}x_2^3(t), & if\ x_2(t) \geq 0, \\ -\frac{1}{6}x_2^3(t), & if\ x_2(t) < 0, \end{cases} \tag{18b}$$

$$x_2(t) - t = 0, \tag{18c}$$

$$x_1(t_0) - p = 0. \tag{18d}$$

Obviously, the index of the DAE system in (18) is $k = 1$ and a unique solution is given by $x_2(t_0) = \dot{x}_1(t_0) = \ddot{x}_1(t_0) = \ddot{x}_2(t_0) = 0$, $x_1(t_0) = p$, $\dot{x}_2(t_0) = 1$.
Hence $z = (x_1(t_0), x_2(t_0), \dot{x}_1(t_0), \dot{x}_2(t_0), \ddot{x}_1(t_0), \ddot{x}_2(t_0))$, we find the (NLP1(p))

$$\min_z \quad x_2^2(t_0), \tag{19a}$$

$$\text{s.t.} \quad \dot{x}_1(t_0) = \begin{cases} \frac{1}{6}x_2^3(t_0), & \text{if } x_2(t_0) \geq 0, \\ -\frac{1}{6}x_2^3(t_0), & \text{if } x_2(t_0) < 0, \end{cases} \tag{19b}$$

$$x_2(t_0) = t_0, \tag{19c}$$

$$\ddot{x}_1(t_0) = \begin{cases} \frac{1}{2}x_2^2(t_0), & \text{if } x_2(t_0) \geq 0, \\ -\frac{1}{2}x_2^2(t_0), & \text{if } x_2(t_0) < 0, \end{cases} \tag{19d}$$

$$\dot{x}_2(t_0) = 1, \tag{19e}$$

$$x_1(t_0) = p. \tag{19f}$$

Since the equation for $\ddot{x}_1(t_0)$ in (19) is not twice continuously differentiable with respect to $x_2(t_0)$, Assumption 3.1 is not fulfilled. Hence Theorem 3.2 and Theorem 3.3 can not be applied. On the other hand, $\ddot{x}_1(t)$ is not used to find consistent initial values for the DAE system (18). Hence we can dispense with Eqs. (19) for $\ddot{x}_1(t_0)$.

4 Weak Solution Differentiability of Consistency Functions

Concerning the difficulties with Example 3.1 in applying Theorem 3.2 and Theorem 3.3 we will show in this section, that the differentiability assumption 3.1 can be appropriately weakened. The idea is to determine relevant equations for finding consistent initial values.

First we reorder the components of the vector $x(t_0)$ of initial values and then partition it as follows:

$$x(t_0) = (x^N(t_0), x^B(t_0))^T$$

with

$$\begin{array}{rcll} x^N(t_0) & = & (x_1^N(t_0), \ldots, x_{n-q}^N(t_0))^T, & x_i^N(t_0) \in \{x_1(t_0), \ldots, x_n(t_0)\}, \\ x^B(t_0) & = & (x_1^B(t_0), \ldots, x_q^B(t_0))^T, & x_i^B(t_0) \in \{x_1(t_0), \ldots, x_n(t_0)\}. \end{array}$$

Here, $x^N(t_0)$ denotes the vector of initial values which can be expressed by the unspecified (free) initial values in the vector $x^B(t_0)$. In the following, we use the notation $\mathbb{N}_{\bar{n}} := \{1, 2, \ldots, \bar{n}\}$, $\bar{n} \in \mathbb{N}$.

Definition 4.1 (Degree of Freedom) Suppose that (7) is solvable. A natural number $q \in \{1, \ldots, n\}$ is called *degree of freedom* in the initial vector $x(t_0)$ of system (7), i.e., the dimension of $x^B(t_0)$, if it is the largest number, such that the following holds: $n_B^E \in \mathbb{N}_{(k+1)n+r}$ is the smallest number of equations, such that $\bar{G} = (\bar{G}_1, \ldots, \bar{G}_{n_B^E})^T$ is a collection of equations of (9), and $\hat{z} = (\hat{z}_1, \ldots, \hat{z}_{n_B^V})^T$ is a collection of variables, such that $\bar{G}(\hat{z}, p) = 0$ can be transformed into

$$\begin{pmatrix} x^N(t_0) \\ \dot{x}(t_0) \end{pmatrix} = \tilde{G}(x^B(t_0), p). \tag{20}$$

Remark 4.1 Herewith, the dependent initial values $x_i^N(t_0)$ are expressed as functions of the free initial values $x_j^B(t_0)$. The second equation in (20) corresponds to the underlying ODE (5) and since (7) is assumed to be solvable, there always exists a transformation like (20).

Definition 4.2 (Basis of Necessary Variables and Equations) Let $q \in \{1, \ldots, n\}$ be the degree of freedom in (7). Let $\bar{G} = (\bar{G}_1, \ldots, \bar{G}_{n_B^E})^T$ and $\hat{z} = (\hat{z}_1, \ldots, \hat{z}_{n_B^V})^T$ be defined as in Definition 4.1. Furthermore let $z^* = (z_1^*, \ldots z_{n_B^E}^*)^T$, $z_i^* \in \{\hat{z}_1, \ldots, \hat{z}_{n_B^V}\}$, $i = 1, \ldots, n_B^E$, with $\bar{G}_{z^*}$ nonsingular. We call the set $B^V \subseteq (\mathbb{N}_{k+2} \cup \{0\}) \times \mathbb{N}_n$ of tuples of indices *basis of necessary variables* for Problem 2.1 and $B^E \subseteq (\mathbb{N}_{k+1} \cup \{0\}) \times \mathbb{N}_n$ of tuples of indices *basis of necessary equations* for Problem 2.1 if

a) $(1, j) \in B^V$, if $x_j(t_0) \in \{x_1^N(t_0), \ldots, x_{n-q}^N(t_0)\}$,
$(2, j) \in B^V$, for all $j \in \mathbb{N}_n$,
$(i, j) \in B^V$, if $x_j^{(i-1)} = \frac{d^{i-1}}{dt^{i-1}} x_j(t_0) \in \{z_1^*, \ldots, z_{n_B^E}^*\}$, $i \geq 3$

b) $(0,j) \in B^E$, if $\varphi_j(t_0) \in \{\bar{G}_1, \ldots, \bar{G}_{n_B^E}\}$,
$(i,j) \in B^E$, if $\frac{d^{i-1}}{dt^{i-1}} F_j(x(t_0), \dot{x}(t_0), p) \in \{\bar{G}_1, \ldots, \bar{G}_{n_B^E}\}$.

Remark 4.2

- In the sets of tuples B^V and B^E, resp., the first index is associated with the derivative and the second with the component of the vector $x(t_0)$, resp., of the equation in (9).
- For a given problem the determination of a basis of necessary variables and a basis of necessary equations is a demanding task, since a carefully analysis of all functions in Problem 2.1 is necessary, compare Pantelides (Ref. 4). The intention of this paper is the analysis of differentiability properties of consistency functions, hence we do not discuss how to obtain these bases.

Those definitions ensure that if a basis of necessary variables and equations exist, the system (9) can be transformed into an ODE system like (5), where the maximal information on $x(t_0)$ is exploited by the consistency equations. Note, that neither the basis of necessary variables nor of equations need to be unique as well as their dimensions. Furthermore the regularity condition in Definition 4.2 ensures the applicability of the implicit function theorem and herewith the following

Theorem 4.1 (Weak Solution Differentiability) Let B^V be a basis of necessary variables of dimension n_B^E, and let B^E be a basis of necessary equations for Problem 2.1. Furthermore let $\hat{z} = (\hat{z}_1, \ldots, \hat{z}_{n_B^E})^T$ be necessary variables B^V and $\bar{G}$ necessary equations B^E. As collection of necessary variables and initial variables $x(t_0)$ we define $C := \{\hat{z}_1, \ldots, \hat{z}_{n_B^E}\} \cup \{x_1(t_0), \ldots, x_n(t_0)\} = \{z_1, \ldots, z_{\hat{n}}\}$, $\hat{n} = n_B^E + q$, and $z := (z_1, \ldots, z_{\hat{n}})^T$. Suppose that (z^0, μ^0) satisfies Assumption 3.1 for the problem

$$\text{(NLP2(p))} \qquad \min_z \quad g(z,p), \tag{21a}$$

$$\text{s.t.} \quad \bar{G}(z,p) = 0. \tag{21b}$$

Then

(a) (z^0, μ^0) is a strong local minimum of (NLP2(p^0)),

(b) there exists a neighborhood $P^0 \subseteq P$ of $p = p^0$ and unique continuously differentiable functions $z : P^0 \longrightarrow \mathbb{R}^{(k+2)n}$, $\mu : P^0 \longrightarrow \mathbb{R}^{(k+1)n+r}$ with the following:

(i) $z(p^0) = z^0$,

(ii) $\mu(p^0) = \mu^0$,

(iii) for all $p \in P^0$: $z(p)$, $\mu(p)$ satisfy Assumption 3.1 for the perturbed problem (NLP2(p)). In particular $(z(p), \mu(p))$ is a unique strong local minimum of (NLP2(p)),

(c) the first order derivatives of $z(p)$ and $\mu(p)$ are given by

$$\begin{pmatrix} \frac{dz}{dp}(p^0) \\ \frac{d\mu}{dp}(p^0) \end{pmatrix} = - \begin{pmatrix} L_{zz}(z^0, \mu^0) & \bar{G}_z(z^0)^T \\ \bar{G}_z(z^0) & 0 \end{pmatrix}^{-1} \begin{pmatrix} L_{zp}(z^0, \mu^0) \\ \bar{G}_p(z^0) \end{pmatrix}, \tag{22}$$

where the Lagrangian L in (22) is given by

$$L(z, \mu, p) := g(z, p) + \mu^T \bar{G}(z, p). \tag{23}$$

The proof of this theorem is equivalent to that of Theorem 3.2 and will not be discussed here. Besides the weaker assumptions, this formulation is especially advantageous in practical applications, e.g., for semi-explicit DAE systems, since the number of variables and equations in (NLP2(p)) is generally (considerably) reduced in comparison to (NLP1(p)).

Again as a direct consequence we find:

Theorem 4.2 (Weak Differentiability Conditions for the Consistency Function con(p)) Suppose there exist consistent initial values for Problem 2.1 and the assumptions in Theorem 4.1 are fulfilled for a reference parameter $p = p^0$. Without loss of generality let $z_i = x_i(t_0)$, $z_{n+i} = \dot{x}_i(t_0)$ for $i = 1, \ldots, n$ and z as in Theorem 4.1. Then, for $x_0(t_0) = (x_1(t_0), \ldots, x_n(t_0))^T$, $\dot{x}_0(t_0) = (\dot{x}_1(t_0), \ldots, \dot{x}_n(t_0))^T$ there exists a neighborhood $P^0 \subseteq P$ of p^0 such that $con(p) := (x_0(p), \dot{x}_0(p))^T$ defines a C^1 consistency function on P^0.

This implies that for p near to p^0 the unperturbed solution (z^0, μ^0) can be embedded into a C^1–family of perturbed optimal solutions $(z(p), \mu(p))$ for (NLP(p)) with $(z(p_0), \mu(p_0)) = (z_0, \mu_0)$. Note again, that in general it is not an easy task to specify a basis of necessary variables and a basis of necessary equations. Hence a carefully analysis of all functions in Problem 2.1 is necessary.

5 Examples

Three different examples will be discussed in this section. In general the nominal perturbation parameter p^0 represents a fixed value. In the sequel we evaluate the theoretical results from the previous sections in an analytical way. This helps us to demonstrate, that the derivatives of the consistency function defined in Theorem 3.3 [resp. Theorem 4.2] calculated by formula (17) [resp. formula (22)] coincide with a direct differentiation of the solutions obtained after evaluating the necessary conditions (13).

Example 5.1 As the first example let us revisit Example 3.1, which could not be solved according to Theorem 3.2 [resp. Theorem 3.3] because of the too strong

differentiability assumptions. By defining the basis of necessary variables and equations

$$B^V := \{(1,1),(1,2),(2,1),(2,2)\}, \tag{24a}$$
$$B^E := \{(0,1),(1,1),(1,2),(2,2)\} \tag{24b}$$

we obtain for $t_0 = 0$:

$$z := (x_1(t_0), x_2(t_0), \dot{x}_1(t_0), \dot{x}_2(t_0))^T, \tag{25a}$$
$$\bar{G}(z,p^0) := \begin{pmatrix} \dot{x}_1(t_0) - \frac{1}{6}x_2(t_0)^3 \\ x_2(t_0) - t_0 \\ \dot{x}_2(t_0) - 1 \\ x_1(t_0) - p^0 \end{pmatrix} = 0, \qquad \text{if } x_2(t_0) \geq 0, \tag{25b}$$
$$\bar{G}(z,p^0) := \begin{pmatrix} \dot{x}_1(t_0) + \frac{1}{6}x_2(t_0)^3 \\ x_2(t_0) - t_0 \\ \dot{x}_2(t_0) - 1 \\ x_1(t_0) - p^0 \end{pmatrix} = 0, \qquad \text{if } x_2(t_0) < 0. \tag{25c}$$

For $\mu \in \mathbb{R}^4$ the Lagrangian is given by $L(z,\mu,p^0) = x_2(t_0)^2 + \mu^T \bar{G}(z,p^0)$. Evaluating the necessary conditions of Theorem 3.1 yields $L_z(z,\mu,p^0) = (x_2(t_0)^2)_z + \mu^T \bar{G}_z(z,p^0) = 0$ and, together with the equality constraints (25), we obtain

$$\bar{z} = (p^0, 0, 0, 1)^T, \tag{26a}$$
$$\bar{\mu} = (0,0,0,0)^T \tag{26b}$$

as a candidate for an optimal solution of

$$\min_z \quad x_2(t_0)^2, \tag{27a}$$
$$\text{s.t.} \quad \bar{G}(z,p^0) = 0. \tag{27b}$$

To check the second order sufficient conditions for $\bar{z}$ and $\bar{\mu}$ we calculate

$$L_{zz}(\bar{z},\bar{\mu},p^0) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \tag{28}$$

and

$$\bar{G}_z(\bar{z},p^0) = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \tag{29}$$

Since $\bar{G}_z(\bar{z},p^0)$ is of full rank we find $\mathrm{Ker}(\bar{G}_z(\bar{z},p^0)) = (0,0,0,0)^T$ and the Hessian $L_{zz}(\bar{z},\bar{\mu},p^0)$ in (28) is positive definite on $\mathrm{Ker}(\bar{G}_z(\bar{z},p^0))\backslash\{0\}$ (compare Assumption 3.1.5).Hence $(\bar{z},\bar{\mu})$ is a strong local minimum of (27). Since all differentiability properties are fulfilled we can apply Theorem 4.1 and Theorem 4.2 and find that there exists a neighborhood $P(p^0)$ for any reference parameter $p = p^0$ and a unique C^1 consistency function $con : P(p^0) \longrightarrow \mathbb{R}^{2n}$:

$$\mathrm{con(p)} := (\mathrm{x_0(p)}, \dot{\mathrm{x}}_0(\mathrm{p}))^{\mathrm{T}} = (\mathrm{z_1(p)}, \mathrm{z_2(p)}, \mathrm{z_3(p)}, \mathrm{z_4(p)})^{\mathrm{T}}. \tag{30}$$

The derivative $\frac{d\text{con}}{dp}(p^0)$ can be calculated with

$$L_{zp}(\bar{z},\bar{\mu},p^0) = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \qquad \bar{G}_p(\bar{z},p^0) = \begin{pmatrix} 0 \\ 0 \\ 0 \\ -1 \end{pmatrix}, \tag{31}$$

and

$$\begin{pmatrix} L_{zz}(\bar{z},\bar{\mu},p^0) & \bar{G}_z(\bar{z},p^0)^T \\ \bar{G}_z(\bar{z},p^0) & 0 \end{pmatrix}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \tag{32}$$

by applying Formula (22) as

$$\frac{d\text{con}}{dp}(p^0) = (1,0,0,0). \tag{33}$$

This coincides with the direct differentiation of $\bar{z}$ in (26) with respect to p^0.

Example 5.2 (Implicit Problem) Assume the following implicitly given problem of index $k = 1$:

$$\begin{pmatrix} x_2(t) & -x_3(t) & 0 \\ -x_2(t) & x_3(t) & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \\ \dot{x}_3(t) \end{pmatrix} - \begin{pmatrix} x_3(t) \\ x_1(t) \\ x_1(t) + x_2(t) \end{pmatrix} = 0. \tag{34}$$

We try to find consistent initial values for $(x_2(t_0), x_3(t_0))$ near to given values (p_1^0, p_2^0):

$$\begin{array}{ll} \min\limits_{\tilde{z}} & (x_2(t_0) - p_1^0)^2 + (x_3(t_0) - p_2^0)^2, \\ \text{subject to} & \text{the DAE system (34) and its first derivatives,} \end{array} \tag{35}$$

where $\tilde{z}$ is defined by

$$\tilde{z} := (x(t_0), \ldots, x^{(k+1)}(t_0))^T. \tag{36}$$

All differentiability assumptions hold for Theorem 3.2 and Theorem 3.3. However we use Theorem 4.1 and Theorem 4.2 and follow the ideas of Section 4. A basis of necessary variables and equations is given by

$$B^V := \{(1,1),(1,2),(2,1),(2,2),(2,3),(3,1)\}, \tag{37a}$$

$$B^E := \{(1,1),(1,2),(1,3),(2,1),(2,2),(2,3)\}. \tag{37b}$$

This gives

$$\begin{aligned} z &:= (x_1(t_0), x_2(t_0), x_3(t_0), \dot{x}_1(t_0), \dot{x}_2(t_0), \dot{x}_3(t_0), \ddot{x}_1(t_0))^T, \\ \bar{G}(z,p^0) &:= \begin{pmatrix} x_2\dot{x}_1 - x_3 - x_3\dot{x}_2 \\ x_3\dot{x}_2 - x_1 - x_2\dot{x}_1 \\ x_2 + x_1 \\ \dot{x}_2\dot{x}_1 + x_2\ddot{x}_1 - \dot{x}_3 - \dot{x}_2\dot{x}_3 - x_3\ddot{x}_2 \\ \dot{x}_3\dot{x}_2 + x_3\ddot{x}_2 - \dot{x}_1 - \dot{x}_1\dot{x}_2 - x_2\ddot{x}_1 \\ \dot{x}_2 + \dot{x}_1 \end{pmatrix} = 0, \end{aligned} \tag{38}$$

where all components in $\bar{G}(z, p^0)$ are evaluated at t_0. Please note, that the component $\ddot{x}_2$ occurs in (38). But, as we will see later, it turns out that this variable has no influence on the initial values $x(t_0)$ and $\dot{x}(t_0)$.

For $\mu \in \mathbb{R}^6$ the Lagrangian is given by $L(z, \mu, p^0) = (x_2(t_0) - p_1^0)^2 + (x_3(t_0) - p_2^0)^2 + \mu^T \bar{G}(z, p^0)$. The equality constraints from $\bar{G}(z, p^0) = 0$ in (38) and the necessary conditions in Theorem 3.1 yield

$$\bar{z} := (-\frac{1}{2}(p_1^0 + p_2^0), \frac{1}{2}(p_1^0 + p_2^0), \frac{1}{2}(p_1^0 + p_2^0), \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \ddot{x}_2,)^T, \tag{39a}$$

$$\bar{\mu} := (p_1^0 - p_2^0, p_1^0 - p_2^0, p_1^0 - p_2^0, 0, 0, 0,)^T. \tag{39b}$$

The Hessian of the Lagrangian evaluated at $(\bar{z}, \bar{\mu})$ is given by

$$L_{zz}(\bar{z}, \bar{\mu}, p^0) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \tag{40}$$

while the Jacobian of the constraints yield for $\beta = \frac{1}{2}(p_1^0 + p_2^0)$,

$$\bar{G}_z(\bar{z}, p^0) = \begin{pmatrix} 0 & \frac{1}{2} & -\frac{1}{2} & \beta & -\beta & 0 & 0 \\ -1 & -\frac{1}{2} & -\frac{1}{2} & -\beta & \beta & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \ddot{x}_2 & -\ddot{x}_2 & -\frac{1}{2} & 1 & -\frac{1}{2} & \beta \\ 0 & -\ddot{x}_2 & \ddot{x}_2 & -\frac{1}{2} & -1 & -\frac{1}{2} & -\beta \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 \end{pmatrix}. \tag{41}$$

The Jacobian $\bar{G}_z(\bar{z}, p^0)$ is of maximal rank, hence the dimension of the kernel is one. We find $\mathrm{Ker}(\bar{G}_z(\bar{z}, p^0)) = \{v \in \mathbb{R}^7 : v = (-\alpha, \alpha, \alpha, 0, 0, 0, 0)^T,\ \alpha \in \mathbb{R}\}$ and thus obtain

$$v^T L_{zz}(\bar{z}, \bar{\mu}, p^0) v = 4\alpha^2 \tag{42}$$

for $v \in \mathrm{Ker}(\bar{G}_z(\bar{z}, p^0))$. Hence, the Hessian is positive definite on $\mathrm{Ker}(\bar{G}_z(\bar{z}, p^0)) \backslash \{0\}$ (compare Assumption 3.1.5). Thus $(\bar{z}, \bar{\mu})$ is a strong local minimum of (34). Since all differentiability properties are fulfilled we can apply Theorem 4.1 and Theorem 4.2 and find that there exists a neighborhood $P(p^0)$ for any reference parameter $p = p^0$ and a unique C^1 consistency function $con : P(p^0) \longrightarrow \mathbb{R}^{2n}$:

$$\mathrm{con(p)} := (\mathrm{x_0(p)}, \dot{\mathrm{x}}_0(\mathrm{p}))^{\mathrm{T}} = (\mathrm{z_1(p)}, \mathrm{z_2(p)}, \ldots, \mathrm{z_6(p)})^{\mathrm{T}}. \tag{43}$$

With

$$L_{zp}(\bar{z}, \bar{\mu}, p^0) = \begin{pmatrix} 0 & 0 \\ -2 & 0 \\ 0 & -2 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \qquad \bar{G}_p(\bar{z}, p^0) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \tag{44}$$

and by formula (22) the derivative $\frac{d\text{con}}{dp}(p^0)$ reads as

$$\frac{d\text{con}}{dp_1}(p_1^0) = \frac{d\text{con}}{dp_2}(p_2^0) = (-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0). \tag{45}$$

This indicates, that small deviations in $p = (p_1, p_2)$ cause the consistent initial values for $x(t_0)$ to change. Especially $x_1(t_0)$ is influenced by this perturbations even though the deviations are only directly coupled with $x_2(t_0)$ and $x_3(t_0)$ in the objective function. The derivative $\frac{d\text{con}}{dp}(p^0)$ coincides with a direct differentiation of the first six arguments of $\bar{z}$ in (39) with respect to p^0.

Example 5.3 (Mathematical Pendulum) The equations of motion of a mathematical pendulum with mass m and length $l = 1$ are given by

$$\begin{aligned}
\dot{x}_1(t) &= x_3(t), && \text{(46a)}\\
\dot{x}_2(t) &= x_4(t), && \text{(46b)}\\
m\dot{x}_3(t) &= -2\,x_5(t)\,x_1(t), && \text{(46c)}\\
m\dot{x}_4(t) &= -mg_0 - 2\,x_5(t)\,x_2(t), && \text{(46d)}\\
0 &= x_1(t)^2 + x_2(t)^2 - 1. && \text{(46e)}
\end{aligned}$$

Here, $x_1(t), \ldots, x_4(t)$ denote the differential variables and $x_5(t)$ denotes the algebraic variable.

Threefold differentiation of the algebraic constraint $0 = x_1(t)^2 + x_2(t)^2 - 1$ w.r.t. time yields the possibility to transform (46) to an ODE system, hence the differentiation index is $k = 3$. Let additional initial values

$$\begin{aligned}
x_1(t_0) &= \frac{1}{\sqrt{2}}, && \text{(47a)}\\
x_2(t_0) &= -\frac{1}{\sqrt{2}} && \text{(47b)}
\end{aligned}$$

be given.
In the sequel we investigate the problem

$$\begin{aligned}
\min_{\hat{z}} \quad & (x_3(t_0) - s)^2 + (x_4(t_0) - s)^2, && \text{(48a)}\\
\text{s.t.} \quad & \text{equation (46), and its derivatives up to order 3, and (47)} && \text{(48b)}
\end{aligned}$$

with $\hat{z}$ defined by

$$\hat{z} := (x(t_0), \ldots, x^{(k+1)}(t_0))^T. \tag{49}$$

As perturbation parameter we choose $p = (m, g_0, s)$. Please note, that similar problems arise in connection with direct solution methods for optimal control problems with DAE systems of higher index and free initial values.

All differentiability assumptions hold for Theorem 3.2 and Theorem 3.3. However, due to the high dimension we favour Theorem 4.1 and Theorem 4.2 and follow

the ideas of Section 4. A basis of necessary variables and equations is given by

$$\begin{aligned} B^V &:= \{(1,1),(1,2),(1,3),(1,4),(2,1),(2,2),(2,3),(2,4),(2,5),\\ &\quad (3,1),(3,2),(3,3),(3,4),(4,1),(4,2)\}, \qquad (50a)\\ B^E &:= \{(0,1),(0,2),(1,1),(1,2),(1,3),(1,4),(2,1),(2,2),(2,3),\\ &\quad (2,4),(2,5),(3,1),(3,2),(3,5),(4,5)\}. \qquad (50b) \end{aligned}$$

For a reference parameter $p^0 = (m^0, g_0^0, s^0)$ this leads to the optimization variables and equations, resp.,

$$\begin{aligned} z &:= (x_1,\ldots,x_5,\dot{x}_1,\ldots,\dot{x}_5,\ddot{x}_1,\ldots,\ddot{x}_4,\dddot{x}_1,\ddddot{x}_2)^T,\\ \bar{G}(z,p^0) &:= \begin{pmatrix} \dot{x}_1 - x_3 \\ \dot{x}_2 - x_4 \\ m^0\dot{x}_3 + 2x_5x_1 \\ m^0\dot{x}_4 + m^0g_0^0 + 2x_5x_2 \\ \ddot{x}_1 - \dot{x}_3 \\ \ddot{x}_2 - \dot{x}_4 \\ m^0\ddot{x}_3 + 2\dot{x}_1x_5 + 2x_1\dot{x}_5 \\ m^0\ddot{x}_4 + 2\dot{x}_2x_5 + 2x_2\dot{x}_5 \\ 2x_1\dot{x}_1 + 2x_2\dot{x}_2 \\ \dddot{x}_1 - \ddot{x}_3 \\ \ddddot{x}_2 - \ddot{x}_4 \\ 2\dot{x}_1^2 + 2x_1\ddot{x}_1 + 2\dot{x}_2^2 + 2x_2\ddot{x}_2 \\ 6\dot{x}_1\ddot{x}_1 + 2x_1\,\dddot{x}_1 + 6\dot{x}_2\ddot{x}_2 + 2x_2\,\ddddot{x}_2 \\ x_1 - \frac{1}{\sqrt{2}} \\ x_2 + \frac{1}{\sqrt{2}} \end{pmatrix} = 0, \end{aligned} \qquad (51)$$

where all components in z and $\bar{G}(z,p^0)$ are evaluated at t_0. For $\mu \in \mathbb{R}^{15}$ the Lagrangian is given by $L(z,\mu,p^0) = (x_3(t_0) - s^0)^2 + (x_4(t_0) - s^0)^2 + \mu^T\bar{G}(z,p^0)$. The equality constraints from $\bar{G}(z,p^0) = 0$ in (51) and the necessary conditions in Theorem 3.1 yield $\bar{\mu} = 0 \in \mathbb{R}^{15}$ and

$$\bar{z} = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \\ s^0 \\ s^0 \\ \frac{1}{4}m^0(\sqrt{2}g_0^0 + 4(s^0)^2) \\ s^0 \\ s^0 \\ -\frac{1}{2}g_0^0 - \sqrt{2}(s^0)^2 \\ -\frac{1}{2}g_0^0 + \sqrt{2}(s^0)^2 \\ -\frac{3}{2}m^0g_0^0s^0 \\ -\frac{1}{2}g_0^0 - \sqrt{2}(s^0)^2 \\ -\frac{1}{2}g_0^0 + \sqrt{2}(s^0)^2 \\ s^0(\sqrt{2}g_0^0 - 2(s^0)^2) \\ -2s^0(\sqrt{2}g_0^0 + (s^0)^2) \\ s^0(\sqrt{2}g_0^0 - 2(s^0)^2) \\ -2s^0(\sqrt{2}g_0^0 + (s^0)^2) \end{pmatrix}. \qquad (52)$$

The Hessian of the Lagrangian is defined by $A := L_{zz}(\bar{z}, \bar{\mu}, m^0, g_0^0, s^0)$ with

$$A = (A)_{i,j} = \begin{cases} 2, & \text{if} \quad i = j = 3, \text{ or } i = j = 4, \\ 0, & \text{else,} \end{cases} \tag{53}$$

while the Jacobian of the constraints yield

$$\bar{G}_z(\bar{z}, p^0) = \begin{pmatrix}
0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
\beta_1 & 0 & 0 & 0 & \frac{\sqrt{2}}{m^0} & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & \beta_1 & 0 & 0 & -\frac{\sqrt{2}}{m^0} & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
-3s^0 g_0^0 & 0 & 0 & 0 & 2\frac{s^0}{m^0} & \beta_1 & 0 & 0 & 0 & \frac{\sqrt{2}}{m^0} & 0 & 0 & 1 & 0 & 0 & 0 \\
0 & -3s^0 g_0^0 & 0 & 0 & 2\frac{s^0}{m^0} & 0 & \beta_1 & 0 & 0 & -\frac{\sqrt{2}}{m^0} & 0 & 0 & 0 & 1 & 0 & 0 \\
2s^0 & 2s^0 & 0 & 0 & 0 & \sqrt{2} & -\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 1 \\
-\beta_2 & \beta_2 & 0 & 0 & 0 & 4s^0 & 4s^0 & 0 & 0 & 0 & \sqrt{2} & -\sqrt{2} & 0 & 0 & 0 & 0 \\
-\beta_3 & 2\beta_3 & 0 & 0 & 0 & -\beta_4 & \beta_4 & 0 & 0 & 0 & 6s^0 & 6s^0 & 0 & 0 & \sqrt{2} & -\sqrt{2} \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix} \tag{54}$$

with

$$\beta_1 = 2(s^0)^2 + \frac{1}{\sqrt{2}} g_0^0, \qquad \beta_2 = 2\sqrt{2}(s^0)^2 - g_0^0, \tag{55a}$$

$$\beta_3 = -2s^0(\sqrt{2} g_0^0 - 2(s^0)^2), \qquad \beta_4 = 6\sqrt{2}(s^0)^2 - 3g_0^0. \tag{55b}$$

The Jacobian is of maximal rank hence the dimension of the kernel is one and given by

$$\begin{aligned} &\mathrm{Ker}(\bar{G}_z(\bar{z}, p^0)) = \{v \in \mathbb{R}^{16} : \\ \text{with} \quad &v = (0, 0, c_2, c_2, c_3, c_2, c_2, -c_4, c_4, c_5, -c_4, c_4, \alpha, c_6, \alpha, c_6)^T, \ \alpha \in \mathbb{R}\} \end{aligned} \tag{56}$$

$$c_1 = g_0^0 - 3\sqrt{2}(s^0)^2, \quad c_2 = \frac{\alpha}{\sqrt{2} c_1}, \qquad c_3 = \frac{\alpha s^0 \sqrt{2} m^0}{c_1}, \tag{57a}$$

$$c_4 = 2\frac{\alpha s^0}{c_1}, \qquad c_5 = -\frac{3\alpha m^0 g_0^0 \sqrt{2}}{4c_1}, \quad c_6 = -\frac{\alpha(2g_0^0 + 3\sqrt{2}(s^0)^2)}{c_1}. \tag{57b}$$

Furthermore we get

$$v^T L_{zz}(\bar{z}, \bar{\mu}, p^0) v = \frac{2\alpha^2}{(g_0^0 - 3\sqrt{2}(s^0)^2)^2}, \tag{58}$$

independent on m^0, hence the Hessian is positive definite on $\mathrm{Ker}(\bar{G}_z(\bar{z}, p^0)) \backslash \{0\}$ (compare Assumption 3.1.5). Thus $(\bar{z}, \bar{\mu})$ is a strong local minimum of (34). Since all differentiability properties are fulfilled we can apply Theorem 4.1 and Theorem 4.2 and find that there exists a neighborhood $P(p^0)$ for any reference parameter $p = p^0$ and a unique C^1 consistency function $\mathrm{con} : \mathrm{P}(\mathrm{p}^0) \longrightarrow \mathbb{R}^{2\mathrm{n}}$:

$$\mathrm{con}(\mathrm{p}) := (\mathrm{x}_0(\mathrm{p}), \dot{\mathrm{x}}_0(\mathrm{p}))^{\mathrm{T}} = (\mathrm{z}_1(\mathrm{p}), \mathrm{z}_2(\mathrm{p}), \ldots, \mathrm{z}_{10}(\mathrm{p}))^{\mathrm{T}}. \tag{59}$$

From $B := L_{zp}(\bar{z}, \bar{\mu}, m^0, g_0^0, s^0)$ with

$$B = (B)_{i,j} = \begin{cases} -2, & \text{if} \quad i = 3,\ j = 3,\ \text{or}\ i = 4,\ j = 3, \\ 0, & \text{else}, \end{cases} \tag{60}$$

and $C := \bar{G}_p(\bar{z}, m^0, g_0^0, s^0)$ with

$$C = (C)_{i,j} = \begin{cases} -\frac{\sqrt{2}g_0^0 + 4(s^0)^2}{2\sqrt{2}m^0}, & \text{if} \quad i = 3,\ j = 1, \\ \frac{\sqrt{2}g_0^0 + 4(s^0)^2}{2\sqrt{2}m^0}, & \text{if} \quad i = 4,\ j = 1, \\ 1, & \text{if} \quad i = 4,\ j = 2, \\ -\frac{s^0(\sqrt{2}g_0^0 + 4(s^0)^2) - 3\sqrt{2}g_0^0 s^0}{2m^0}, & \text{if} \quad i = 7,\ j = 1, \\ -\frac{s^0(\sqrt{2}g_0^0 + 4(s^0)^2) + 3\sqrt{2}g_0^0 s^0}{2m^0}, & \text{if} \quad i = 8,\ j = 1, \\ 0, & \text{else}, \end{cases} \tag{61}$$

the derivative $\frac{d\text{con}}{dp}(p^0)$ reads as

$$\frac{d\text{con}}{dp}(p^0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & - \\ \frac{1}{4}\sqrt{2}(2\sqrt{2}(s^0)^2 + g_0^0) & -\frac{1}{4}\sqrt{2}m^0 & 2m^0 s \\ 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & -\frac{1}{2} & -2\sqrt{2}s^0 \\ 0 & -\frac{1}{2} & 2\sqrt{2}s^0 \\ -\frac{3}{2}s^0 g_0^0 & -\frac{3}{2}m^0 s^0 & -\frac{3}{2}g_0^0 m^0 \end{pmatrix}. \tag{62}$$

Again the sensitivity derivative coincides with a direct differentiation of the first ten arguments of $\bar{z}$ in (52) with respect to (m^0, g^0, s^0).

6 Conclusions

In this article parametric differential–algebraic systems are considered. Due to the parameter dependency of the DAE system, in general the solution and in particular consistent initial values depend on these parameters. Therefore the concept of consistency functions is introduced. A consistency function maps a given parameter to a consistent initial value for the DAE system. This consistent initial value need not to be unique, such that the resulting degrees of freedom in the choice of the consistent value can be exploited to optimize a given performance index. This leads to optimal consistent initial values, which play an important role in conjunction with direct shooting methods for the numerical solution of optimal control problems with DAE systems of higher index and free initial values, see Refs. 8,12. One type of such consistency functions is given in Section 2. Of particular importance, e.g. for DAE optimal control problems, is the investigation, under which conditions consistency functions are differentiable functions with respect to the parameters. Therefore necessary and sufficient conditions are stated. It turns out that these conditions are

too restrictive in some cases. An example is given in Section 3. Another class of consistency functions given in Section 4 is based on the idea to extract only relevant information and allows to formulate weaker conditions to obtain the desired differentiability properties. The discussion of several illustrative examples shows the capability of our investigations.

Because the required derivatives of the DAE system can not be provided nor computed analytically, in realistic technical applications it is often not possible to construct a consistency function in the depicted way. Therefore, methods for the *numerical approximation* of consistency functions are needed. A first approach can be found in Büskens and Gerdts (Ref. 8). As mentioned before, differentiable consistency functions can be incorporated in direct shooting techniques for the numerical solution of DAE optimal control problems. The sensitivities, i.e. the differentials of the consistency function w.r.t. parameters, play an import role in the calculation of gradients and Jacobians, e.g. by means of the so called sensitivity DAE system, see Gerdts (Ref. 12).

References

1. Petzold, L.R., *Differential/Algebraic Equations are not ODE's,* SIAM Journal on Scientific and Statistical Computing, Vol. 3, pp. 367–384, 1982.

2. Brenan, K. E., Campbell, S. L., and Petzold, L. R., *Numerical Solution of Initial-Value Problems in Differential-Algebraic Equations*, Classics in Applied Mathematics, SIAM, Philadelphia, Pennsylvania, Vol. 14, 1996.

3. Leimkuhler, B., Petzold, L.R., and Gear, C.W., *Approximation Methods for the Consistent Initialization of Differential-Algebraic Equations,* SIAM Journal on Numerical Analysis, Vol. 28, pp. 205-226, 1991.

4. Pantelides, C.C., *The Consistent Initialization of Differential-Algebraic Equations,* SIAM Journal on Scientific and Statistical Computing, Vol. 9, pp. 213-231, 1988.

5. Lamour, R., *A Shooting Method for Fully Implicit Index-2 Differential-Algebraic Equations,* SIAM Journal on Scientific Computing, Vol. 18, pp. 94-114, 1997.

6. Campbell, S.L., *Least Squares Completions for Nonlinear Differential Algebraic Equations,* Numerische Mathematik, Vol. 65, pp. 77-94, 1993.

7. Campbell, S.L., Kelley, C.T., and Yeomans, K.D., *Consistent Initial Conditions for Unstructured Higher Index DAEs: A Computational Study,* Proceedings Computational Engineering in Systems Applications, Lille, France, pp. 416-421, 1996.

8. Büskens, C., and Gerdts, M., *Numerical Solution of Optimal Control Problems with DAE Systems of Higher Index*, Optimalsteuerungsprobleme in der Luft- und Raumfahrt, SFB 255: Transatmosphärische Flugsysteme, Technische Universität München, München, Germany, pp. 27–38, 2000.

9. Büskens, C., and Gerdts, M., *Computation of Consistent Initial Values for Optimal Control Problems with DAE Systems of Higher Index*, ZAMM, Vol. 81, Supplement 2, pp. 249-250, 2001.

10. Fiacco, A.V., *Introduction to Sensitivity and Stability Analysis in Nonlinear Programming*, Mathematics in Science and Engineering, Academic Press, New York, Vol. 165, 1983.

11. Büskens, C., *Optimization Methods and Sensitivity Analysis for Optimal Control Problems with Control and State Constraints*, PhD Thesis, Institute for Numerical and Instrumental Mathematics, University of Münster, Münster, Germany, 1998 (in German).

12. Gerdts, M., *Numerical Methods for Optimal Control Problems with Differential-Algebraic Equation Systems of Higher Index and their Application in Vehicle Simulation and Mechanics*, Bayreuther Mathematische Schriften, Bayreuth, Germany, Vol. 61, 2001 (in German).

13. Gear, C.W., *The Simultaneous Numerical Solution of Differential-Algebraic Equations,* IEEE Transactions on Circuit Theory, Vol. 18, pp. 89–95, 1971.

14. Campbell, S.L., and Gear, C.W., *The Index of General Nonlinear DAEs,* Numerische Mathematik, Vol. 72, pp. 173–196, 1995.

15. Fletcher, R., *Practical Methods of Optimization*, 2nd Edition, J. Wiley and Sons, Chichester, England, 1997.

Berichte aus der Technomathematik ISSN 1435-7968

http://www.math.uni-bremen.de/zetem/berichte.html

— Vertrieb durch den Autor —

Reports **Stand: 12. August 2004**

98–01. Peter Benner, Heike Faßbender:
An Implicitly Restarted Symplectic Lanczos Method for the Symplectic Eigenvalue Problem, Juli 1998.

98–02. Heike Faßbender:
Sliding Window Schemes for Discrete Least-Squares Approximation by Trigonometric Polynomials, Juli 1998.

98–03. Peter Benner, Maribel Castillo, Enrique S. Quintana-Ortí:
Parallel Partial Stabilizing Algorithms for Large Linear Control Systems, Juli 1998.

98–04. Peter Benner:
Computational Methods for Linear–Quadratic Optimization, August 1998.

98–05. Peter Benner, Ralph Byers, Enrique S. Quintana-Ortí, Gregorio Quintana-Ortí:
Solving Algebraic Riccati Equations on Parallel Computers Using Newton's Method with Exact Line Search, August 1998.

98–06. Lars Grüne, Fabian Wirth:
On the rate of convergence of infinite horizon discounted optimal value functions, November 1998.

98–07. Peter Benner, Volker Mehrmann, Hongguo Xu:
A Note on the Numerical Solution of Complex Hamiltonian and Skew-Hamiltonian Eigenvalue Problems, November 1998.

98–08. Eberhard Bänsch, Burkhard Höhn:
Numerical simulation of a silicon floating zone with a free capillary surface, Dezember 1998.

99–01. Heike Faßbender:
The Parameterized SR Algorithm for Symplectic (Butterfly) Matrices, Februar 1999.

99–02. Heike Faßbender:
Error Analysis of the symplectic Lanczos Method for the symplectic Eigenvalue Problem, März 1999.

99–03. Eberhard Bänsch, Alfred Schmidt:
Simulation of dendritic crystal growth with thermal convection, März 1999.

99–04. Eberhard Bänsch:
Finite element discretization of the Navier-Stokes equations with a free capillary surface, März 1999.

99–05. Peter Benner:
Mathematik in der Berufspraxis, Juli 1999.

99–06. Andrew D.B. Paice, Fabian R. Wirth:
Robustness of nonlinear systems and their domains of attraction, August 1999.

99–07. Peter Benner, Enrique S. Quintana-Ortí, Gregorio Quintana-Ortí:
Balanced Truncation Model Reduction of Large-Scale Dense Systems on Parallel Computers, September 1999.

99–08. Ronald Stöver:
Collocation methods for solving linear differential-algebraic boundary value problems, September 1999.

99–09. Huseyin Akcay:
Modelling with Orthonormal Basis Functions, September 1999.

99–10. Heike Faßbender, D. Steven Mackey, Niloufer Mackey:
Hamilton and Jacobi come full circle: Jacobi algorithms for structured Hamiltonian eigenproblems, Oktober 1999.

99–11. Peter Benner, Vincente Hernández, Antonio Pastor:
On the Kleinman Iteration for Nonstabilizable System, Oktober 1999.

99–12. Peter Benner, Heike Faßbender:
A Hybrid Method for the Numerical Solution of Discrete-Time Algebraic Riccati Equations, November 1999.

99–13. Peter Benner, Enrique S. Quintana-Ortí, Gregorio Quintana-Ortí:
Numerical Solution of Schur Stable Linear Matrix Equations on Multicomputers, November 1999.

99–14. Eberhard Bänsch, Karol Mikula:
Adaptivity in 3D Image Processing, Dezember 1999.

00–01. Peter Benner, Volker Mehrmann, Hongguo Xu:
Perturbation Analysis for the Eigenvalue Problem of a Formal Product of Matrices, Januar 2000.

00–02. Ziping Huang:
Finite Element Method for Mixed Problems with Penalty, Januar 2000.

00–03. Gianfrancesco Martinico:
Recursive mesh refinement in 3D, Februar 2000.

00–04. Eberhard Bänsch, Christoph Egbers, Oliver Meincke, Nicoleta Scurtu:
Taylor-Couette System with Asymmetric Boundary Conditions, Februar 2000.

00–05. Peter Benner:
Symplectic Balancing of Hamiltonian Matrices, Februar 2000.

00–06. Fabio Camilli, Lars Grüne, Fabian Wirth:
A regularization of Zubov's equation for robust domains of attraction, März 2000.

00–07. Michael Wolff, Eberhard Bänsch, Michael Böhm, Dominic Davis:
Modellierung der Abkühlung von Stahlbrammen, März 2000.

00–08. Stephan Dahlke, Peter Maaß, Gerd Teschke:
Interpolating Scaling Functions with Duals, April 2000.

00–09. Jochen Behrens, Fabian Wirth:
A globalization procedure for locally stabilizing controllers, Mai 2000.

00–10. Peter Maaß, Gerd Teschke, Werner Willmann, Günter Wollmann:
Detection and Classification of Material Attributes – A Practical Application of Wavelet Analysis, Mai 2000.

00–11. Stefan Boschert, Alfred Schmidt, Kunibert G. Siebert, Eberhard Bänsch, Klaus-Werner Benz, Gerhard Dziuk, Thomas Kaiser:
Simulation of Industrial Crystal Growth by the Vertical Bridgman Method, Mai 2000.

00–12. Volker Lehmann, Gerd Teschke:
Wavelet Based Methods for Improved Wind Profiler Signal Processing, Mai 2000.

00–13. Stephan Dahlke, Peter Maass:
A Note on Interpolating Scaling Functions, August 2000.

00–14. Ronny Ramlau, Rolf Clackdoyle, Frédéric Noo, Girish Bal:
Accurate Attenuation Correction in SPECT Imaging using Optimization of Bilinear Functions and Assuming an Unknown Spatially-Varying Attenuation Distribution, September 2000.

00–15. Peter Kunkel, Ronald Stöver:
Symmetric collocation methods for linear differential-algebraic boundary value problems, September 2000.

00–16. Fabian Wirth:
The generalized spectral radius and extremal norms, Oktober 2000.

00–17. Frank Stenger, Ahmad Reza Naghsh-Nilchi, Jenny Niebsch, Ronny Ramlau:
A unified approach to the approximate solution of PDE, November 2000.

00–18. Peter Benner, Enrique S. Quintana-Ortí, Gregorio Quintana-Ortí:
Parallel algorithms for model reduction of discrete–time systems, Dezember 2000.

00–19. Ronny Ramlau:
A steepest descent algorithm for the global minimization of Tikhonov–Phillips functional, Dezember 2000.

01–01. Efficient methods in hyperthermia treatment planning:
Torsten Köhler, Peter Maass, Peter Wust, Martin Seebass, Januar 2001.

01–02. Parallel Algorithms for LQ Optimal Control of Discrete-Time Periodic Linear Systems:
Peter Benner, Ralph Byers, Rafael Mayo, Enrique S. Quintana-Ortí, Vicente Hernández, Februar 2001.

01–03. Peter Benner, Enrique S. Quintana-Ortí, Gregorio Quintana-Ortí:
Efficient Numerical Algorithms for Balanced Stochastic Truncation, März 2001.

01–04. Peter Benner, Maribel Castillo, Enrique S. Quintana-Ortí:
Partial Stabilization of Large-Scale Discrete-Time Linear Control Systems, März 2001.

01–05. Stephan Dahlke:
Besov Regularity for Edge Singularities in Polyhedral Domains, Mai 2001.

01–06. Fabian Wirth:
A linearization principle for robustness with respect to time-varying perturbations, Mai 2001.

01–07. Stephan Dahlke, Wolfgang Dahmen, Karsten Urban:
Adaptive Wavelet Methods for Saddle Point Problems - Optimal Convergence Rates, Juli 2001.

01–08. Ronny Ramlau:
Morozov's Discrepancy Principle for Tikhonov regularization of nonlinear operators, Juli 2001.

01–09. Michael Wolff:
Einführung des Drucks für die instationären Stokes–Gleichungen mittels der Methode von Kaplan, Juli 2001.

01–10. Stephan Dahlke, Peter Maaß, Gerd Teschke:
Reconstruction of Reflectivity Desities by Wavelet Transforms, August 2001.

01–11. Stephan Dahlke:
Besov Regularity for the Neumann Problem, August 2001.

01–12. Bernard Haasdonk, Mario Ohlberger, Martin Rumpf, Alfred Schmidt, Kunibert G. Siebert:
h–p–Multiresolution Visualization of Adaptive Finite Element Simulations, Oktober 2001.

01–13. Stephan Dahlke, Gabriele Steidl, Gerd Teschke:
Coorbit Spaces and Banach Frames on Homogeneous Spaces with Applications to Analyzing Functions on Spheres, August 2001.

02–01. Michael Wolff, Michael Böhm:
Zur Modellierung der Thermoelasto-Plastizität mit Phasenumwandlungen bei Stählen sowie der Umwandlungsplastizität, Februar 2002.

02–02. Stephan Dahlke, Peter Maaß:
An Outline of Adaptive Wavelet Galerkin Methods for Tikhonov Regularization of Inverse Parabolic Problems, April 2002.

02–03. Alfred Schmidt:
A Multi-Mesh Finite Element Method for Phase Field Simulations, April 2002.

02–04. Sergey N. Dachkovski, Michael Böhm:
A Note on Finite Thermoplasticity with Phase Changes, Juli 2002.

02–05. Michael Wolff, Michael Böhm:
Phasenumwandlungen und Umwandlungsplastizität bei Stählen im Konzept der Thermoelasto-Plastizität, Juli 2002.

02–06. Gerd Teschke:
Construction of Generalized Uncertainty Principles and Wavelets in Anisotropic Sobolev Spaces, August 2002.

02–07. Ronny Ramlau:
TIGRA – an iterative algorithm for regularizing nonlinear ill–posed problems, August 2002.

02–08. Michael Lukaschewitsch, Peter Maaß, Michael Pidcock:
Tikhonov regularization for Electrical Impedance Tomography on unbounded domains, Oktober 2002.

02–09. Volker Dicken, Peter Maaß, Ingo Menz, Jenny Niebsch, Ronny Ramlau:
Inverse Unwuchtidentifikation an Flugtriebwerken mit Quetschöldämpfern, Oktober 2002.

02–10. Torsten Köhler, Peter Maaß, Jan Kalden:
Time-series forecasting for total volume data and charge back data, November 2002.

02–11. Angelika Bunse-Gerstner:
A Short Introduction to Iterative Methods for Large Linear Systems, November 2002.

02–12. Peter Kunkel, Volker Mehrmann, Ronald Stöver:
Symmetric Collocation for Unstructured Nonlinear Differential-Algebraic Equations of Arbitrary Index, November 2002.

02–13. Michael Wolff:
Ringvorlesung: Distortion Engineering 2
Kontinuumsmechanische Modellierung des Materialverhaltens von Stahl unter Berücksichtigung von Phasenumwandlungen, Dezember 2002.

02–14. Michael Böhm, Martin Hunkel, Alfred Schmidt, Michael Wolff:
Evaluation of various phase-transition models for 100Cr6 for application in commercial FEM programs, Dezember 2002.

03–01. Michael Wolff, Michael Böhm, Serguei Dachkovski:
Volumenanteile versus Massenanteile - der Dilatometerversuch aus der Sicht der Kontinuumsmechanik, Januar 2003.

03–02. Daniel Kessler, Ricardo H. Nochetto, Alfred Schmidt:
A posteriori error control for the Allen-Cahn Problem: circumventing Gronwall's inequality, März 2003.

03–03. Michael Böhm, Jörg Kropp, Adrian Muntean:
On a Prediction Model for Concrete Carbonation based on Moving Interfaces - Interface concentrated Reactions, April 2003.

03–04. Michael Böhm, Jörg Kropp, Adrian Muntean:
A Two-Reaction-Zones Moving-Interface Model for Predicting $Ca(OH)_2$ Carbonation in Concrete, April 2003.

03–05. Vladimir L. Kharitonov, Diederich Hinrichsen:
Exponential estimates for time delay systems, May 2003.

03–06. Michael Wolff, Michael Böhm, Serguei Dachkovski, Günther Löwisch:
Zur makroskopischen Modellierung von spannungsabhängigem Umwandlungsverhalten und Umwandlungsplastizität bei Stählen und ihrer experimentellen Untersuchung in einfachen Versuchen, Juli 2003.

03–07. Serguei Dachkovski, Michael Böhm, Alfred Schmidt, Michael Wolff:
Comparison of several kinetic equations for pearlite transformation in 100Cr6 steel,
Juli 2003.

03–08. Volker Dicken, Peter Maass, Ingo Menz, Jenny Niebsch, Ronny Ramlau:
Nonlinear Inverse Unbalance Reconstruction in Rotor dynamics,
Juli 2003.

03–09. Michael Böhm, Serguei Dachkovski, Martin Hunkel, Thomas Lübben, Michael Wolff:
Übersicht über einige makroskopische Modelle für Phasenumwandlungen im Stahl,
Juli 2003.

03–10. Michael Wolff, Friedhelm Frerichs, Bettina Suhr:
Vorstudie für einen Bauteilversuch zur Umwandlungsplastizität bei der perlitischen Umwandlung des Stahls 100 Cr6,
August 2003.

03–11. Michael Wolff, Bettina Suhr:
Zum Vergleich von Massen- und Volumenanteilen bei der perlitischen Umwandlung der Stähle 100Cr6 und C80,
September 2003.

03–12. Rike Grotmaack, Adrian Muntean:
Stabilitätsanalyse eines Moving-Boundary-Modells der beschleunigten Karbonatisierung von Portlandzementen,
September 2003.

03–13. Alfred Schmidt, Michael Wolff, Michael Böhm:
Numerische Untersuchungen für ein Modell des Materialverhaltens mit Umwandlungsplastizität und Phasenumwandlungen beim Stahl 100Cr6 (Teil 1),
September 2003.

04–01. Liliana Cruz Martin, Gerd Teschke:
A new method to reconstruct radar reflectivities and Doppler information,
Januar 2004.

04–02. Ingrid Daubechies, Gerd Teschke:
Wavelet based image decomposition by variational functionals,
Januar 2004.

04–03. N. Guglielmi, F. Wirth, M. Zennaro:
Complex polytope extremality results for families of matrices,
März 2004.

04–04. I. Daubechies, G. Teschke:
Variational image restoration by means of wavelets: simultaneous decomposition, deblurring and denoising,
April 2004.

04–05. V.L. Kharitonov, E. Plischke:
Lyapunov matrices for time-delay systems,
April 2004.

04–06. Ronny Ramlau:
On the use of fixed point iterations for the regularization of nonlinear ill-posed problems,
Juni 2004.

04–07. Christof Büskens, Matthias Knauer:
Higher Order Real-Time Approximations In Optimal Control of Multibody-Systems For Industrial Robots,
August 2004.

04–08. Christof Büskens, Roland Griesse:
Computational Parametric Sensitivity Analysis of Perturbed PDE Optimal Control Problems with State and Control Constraints,
August 2004.

04–09. Christof Büskens:
Higher Order Real-Time Approximations of Perturbed Control Constrainted PDE Optimal Control Problems with State and Control Constraints,
August 2004.

04–10. Christof Büskens, Matthias Gerdts:
Differentiability of Consistency Functions,
August 2004.